

REPRINTED FROM

**UTILITAS  
MATHEMATICA**

WINNIPEG, CANADA

**Some Remarks on the Interval of Existence of Solutions  
to First Order Initial Value Problems**

**M. Frigon**  
Department of Mathematics  
University of Montreal  
Montreal, CANADA

**D. O'Regan**  
Department of Mathematics and Applied Statistics  
University of Idaho  
Moscow, Idaho 83843  
U.S.A.

**Abstract.** In this paper we obtain "weak solutions" via Topological Transversality to nonlinear first order initial value problems of the form  $y' = f(t; y)$ ,  $t \in [0, T]$ , with  $y(0) = r$ , where  $f: [0, T] \times R^n \rightarrow R^n$  satisfies the Caratheodory Conditions. Our analysis, which is based on the notions of an essential map and on a priori bounds on solutions, is used to examine the dependence of the interval of existence for the above stated problem on its initial data and the nonlinearity in the differential equation.

**1. Introduction.**

Many physical situations give rise to initial value problems of the form

$$\begin{cases} y' = f(t; y), & t \in [0, T] \\ y(0) = r \end{cases} \quad (1.1)$$

where in fact  $f: [0, T] \times R^n \rightarrow R^n$  is discontinuous. In this paper we study the case where  $f$  satisfies the *Caratheodory Conditions*, that is,

- (a) for fixed  $y = (y_1, \dots, y_n) \in R^n$ ,  $f(\cdot; y)$  is measurable on  $[0, T]$ ;
- (b) for all  $t \in [0, T]$ ,  $f(t; \cdot)$  is continuous on  $R^n$ .

Let  $R^n$  denote Euclidean  $n$ -space and  $|v|$  the Euclidean norm. Also let  $L^2((0, T); R^n)$  denote the Banach space of Lebesgue measurable functions  $g$  on  $(0, T)$  which are square integrable with the norm given by  $\|g\|_{L^2} = \left( \int_0^T |g(t)|^2 dt \right)^{\frac{1}{2}}$ . Finally,  $L^\infty((0, T); R^n)$  denotes the space of essentially bounded measurable functions on  $(0, T)$ .

By a weak solution to (1.1) we mean a function  $y$  which is absolutely continuous on  $[0, T]$  with  $y' \in L^2((0, T); R^n)$ ,  $y(0) = r$  and  $y' = f(t; y)$  almost everywhere on  $[0, T]$ . We shall establish under reasonable physical assumptions on  $f$ , that (1.1) has bounded weak solutions. This paper in fact extends results of Lee and O'Regan [12] which deals with the case where  $f$  is continuous. Our analysis is based on the Topological Transversality theorem and known results on Sobolev spaces.

**Preliminary Notation.**

Let  $H^1((0, T); \mathbb{R}^n)$  denote the space of functions  $\mathbf{u}$  which are absolutely continuous on  $[0, T]$  and whose derivative  $\mathbf{u}'$  (which exists almost everywhere) is an element of  $L^2((0, T); \mathbb{R}^n) \cdot H^1((0, T); \mathbb{R}^n)$  with norm.

$$\|\mathbf{u}\|_{H^1} = \|\mathbf{u}\|_{L^2} + \|\mathbf{u}'\|_{L^2}$$

is a Banach space. Also we let

$$H_B^1((0, T); \mathbb{R}^n) = \{\mathbf{u} \in H^1((0, T); \mathbb{R}^n) : \mathbf{u}(0) = \mathbf{r}\}.$$

**Theorem 1.1 (Sobolev Imbedding Theorem).**  $H^1((0, T); \mathbb{R}^n)$  is compactly imbedded into  $C([0, T]; \mathbb{R}^n)$  that is, the imbedding operator  $j: H^1((0, T); \mathbb{R}^n) \rightarrow C([0, T]; \mathbb{R}^n)$  is continuous and completely continuous.

**Theorem 1.2 (Change of variable, [16] or [17]).** Let  $g$  be an absolutely continuous function on  $[a, b]$  with  $g' \in L(a, b)$  and  $g(a) < g(b)$ . If  $f$  is a measurable bounded function on  $[g(a), g(b)]$  then

$$\int_{g(a)}^{g(b)} f(y) dy = \int_a^b f(g(t))g'(t) dt.$$

Proof: (Sketch) Let  $W^{1,1}(a, b)$  denote the space of absolutely continuous functions whose derivative is an element of  $L^1((a, b); \mathbb{R}^n) \equiv L^1(a, b)$ . Put  $G(y) = \int_{g(a)}^y f(x) dx$ , so  $G' = f$  and  $G \in W^{1,1}(g(a), g(b))$ . Then there exists a sequence  $G_n$  such that  $G_n|_{[g(a), g(b)]} \rightarrow G$  in  $W^{1,1}$  and  $\|G_n\|_{L^\infty} \leq M_1, \|G'_n\|_{L^\infty} \leq M_2$ . Also,  $G_n \circ g \in W^{1,1}(a, b)$ ,  $(G_n \circ g)' = (G'_n \circ g)g'$  so  $G_n(g(b)) - G_n(g(a)) = \int_a^b G'_n(g(t))g'(t) dt$ . Now since  $G_n(g(b)) - G_n(g(a)) \rightarrow G(g(b)) - G(g(a)) = \int_{g(a)}^{g(b)} f(x) dx$ , it remains to show

$$\int_a^b G'_n(g(t))g'(t) dt \rightarrow \int_a^b f(g(t))g'(t) dt.$$

One has  $G'_n|_{[g(a), g(b)]} \rightarrow f$  in  $L^1(g(a), g(b))$  so there exists a subsequence  $(G'_{n_k})$  such that  $G'_{n_k}(x) \rightarrow f(x)$  a.e. for  $x \in [g(a), g(b)]$ . Put  $E = \{t \in [a, b] : g'(t) \text{ exists and } g'(t) \neq 0\}$ ,  $Z_0 = \{x \in [g(a), g(b)] : G'_{n_k}(x) \rightarrow f(x)\}$  and  $Z = [g(a), g(b)]|Z_0$ . Let  $B \subset E$  such that  $g(B) \subset Z$ , then  $\text{mes}[g(B)] = 0$  and a little argument yields  $\text{mes}[B] = 0$ . Thus,

$$G'_{n_k}(g(t))g'(t) \rightarrow f(g(t))g'(t) \text{ a.e. } t \in [a, b],$$

with

$$|G'_{n_k}(g(t))g'(t)| \leq M_2|g'(t)| \in L^1(a, b).$$

The result follows from the Lebesgue dominated convergence theorem. ■

## 2. Weak solutions to first order initial value problems.

We begin by considering the problems

$$\begin{cases} y' = f(t; y), & t \in [0, T] \\ y(0) = 0 \end{cases} \quad (2.1)$$

where  $f: [0, T] \times \mathbb{R}^n \rightarrow \mathbb{R}^n$  is a function which satisfies the following hypothesis:

$$f \text{ satisfied the Caratheodory Conditions.} \quad (2.2)$$

$$\text{There exists a function } \psi: [0, \infty) \rightarrow (0, \infty); \psi, \frac{1}{\psi} \in L_{loc}^\infty [0, \infty) \quad (2.3)$$

such that  $|f(t; y)| \leq \psi(|y|)$  for  $t \in [0, T]$ .

**Proposition 2.1.** *Suppose  $f$  satisfies (2.2) and (2.3). Then  $F: C([0, T]; \mathbb{R}^n) \rightarrow L^2((0, T); \mathbb{R}^n)$  given by  $(Fu)(t) = f(t; u(t))$  is defined and continuous.*

Proof: Let  $\epsilon > 0$  be given and suppose  $u_0 \in C([0, T]; \mathbb{R}^n)$ . Consider

$$G_{vm} = \left\{ t \in [0, T]: |v - u_0(t)| < \frac{1}{m} \Rightarrow |f(t; v) - f(t; u_0(t))| < \frac{\epsilon}{L} \right\}$$

where  $v \in \mathbb{R}^n$  and  $L$  is a predetermined constant which will be described below.  $G_{vm}$  is measurable since  $f$  satisfies (2.2). Let  $E_{m\epsilon} = \bigcap_{v \in \mathbb{R}^n} G_{vm}$ . Now  $E_{m\epsilon}$  is measurable and  $E_{1\epsilon} \subset E_{2\epsilon} \subset \dots$ . Also  $\bigcup_{m=1}^\infty E_{m\epsilon} = (0, T)$  for if  $t_0 \in (0, T)$ , then there exists  $m$  such that  $|v - u_0(t_0)| < \frac{1}{m}$  and hence  $|f(t_0; v) - f(t_0; u_0(t_0))| < \frac{\epsilon}{L}$  since  $f(t_0; \cdot)$  satisfies (2.2) and so  $t_0 \in E_{m_0\epsilon}$ . Hence, there exists  $m_0 \in \mathbb{N}$  such that  $\text{mes}(E_{m_0\epsilon}) > T - \frac{\epsilon}{L}$ . Finally, let  $K = \text{ess sup}_{x \leq 1 + \|u_0\|_\infty} \psi(x) < \infty$  and  $\eta < \frac{\epsilon^2}{8K^2}$  where  $\|u_0\|_\infty = \sup_{t \in [0, T]} |u_0(t)|$ . Put  $0 < \delta < \frac{1}{m_0}$  and  $\max\{\frac{\epsilon}{\eta}, (2T)^{\frac{1}{2}}\} < L$ . Let  $u \in C([0, T]; \mathbb{R}^n)$  such that  $\|u - u_0\|_\infty < \delta$ . We will now show that  $\|Fu - Fu_0\|_{L^2} < \epsilon$ . If  $t_0 \in E_{m_0\epsilon}$ , then  $|f(t; u(t)) - f(t; u_0(t))| < \frac{\epsilon}{L}$  and so  $\int_{E_{m_0\epsilon}} |f(t; u(t)) - f(t; u_0(t))|^2 dt < \frac{\epsilon^2 T}{L^2} < \frac{\epsilon^2 T}{2T} = \frac{\epsilon^2}{2}$ . However,  $\text{mes}(E_{m_0\epsilon}^c) < \frac{\epsilon}{L} < \frac{\eta}{\epsilon} = \eta$  and so

$$\begin{aligned} \int_{E_{m_0\epsilon}^c} |f(t; u(t)) - f(t; u_0(t))|^2 dt &\leq 2 \int_{E_{m_0\epsilon}^c} \{[\psi(|u(t)|)]^2 + [\psi(|u_0(t)|)]^2\} dt \\ &\leq 4K^2 \eta < \frac{4K^2 \epsilon^2}{8K^2} = \frac{\epsilon^2}{2}. \end{aligned}$$

Hence  $\|Fu - Fu_0\|_{L^2} < \epsilon$ , so  $F: C([0, T]; \mathbb{R}^n) \rightarrow L^2((0, T); \mathbb{R}^n)$  is continuous. ■

The Sobolev Imbedding Theorem together with Proposition 2.1 are now used to extend Theorem 2.1 of [8] for the new class of problems (2.1).

**Theorem 2.2.** Let  $f: [0, T] \times R^n \rightarrow R^n$  satisfy (2.2), (2.3) and  $0 \leq \lambda \leq 1$ . Suppose there is a constant  $K$  independent of  $\lambda$  such that  $\|y\|_{H^1} \leq K$  for each solution  $y(t)$  to

$$\begin{cases} y' = \lambda f(t; y), & t \in (0, T) \\ y(0) = 0. \end{cases} \quad (2.1)_\lambda$$

Then (2.1) has a solution  $y$  in  $H^1((0, T); R^n)$ .

Proof: Let  $B$  denote the set of functions which satisfy the initial condition  $y(0) = 0$ . Let  $\bar{V} = \{u \in H_B^1((0, T); R^n) : \|y\|_{H^1} \leq K + 1\}$  and define  $F_\lambda: C([0, T]; R^n) \rightarrow L^2((0, T); R^n)$  by  $(F_\lambda v)(t) = \lambda f(t; v(t))$ . Now  $F_\lambda$  is continuous by Proposition 2.1. We also have the imbedding  $j: H_B^1((0, T); R^n) \rightarrow C([0, T]; R^n)$  defined by  $ju = u$  completely continuous by Theorem 1.1. Finally, we define  $N: H_B^1((0, T); R^n) \rightarrow L^2((0, T); R^n)$  by  $Ny = y'$ . Clearly,  $N$  is linear and continuous.  $N$  is also one-to-one since if  $Ny = 0$  the absolute continuity of  $y$  together with the initial condition yields  $y \equiv 0$ . To show  $N$  is onto let  $f(t) \in L^2((0, T); R^n)$  and take  $y(t) = \int_0^t f(u) du$ . Clearly,  $y$  is absolutely continuous,  $y(0) = 0$ ,  $y' = f(t)$  almost everywhere and  $y' \in L^2((0, T); R^n)$ . It follows from Theorem 5.10 of [14] that  $N^{-1}$  is a bounded linear operator. Thus,  $H_\lambda = N^{-1}F_\lambda j: \bar{V} \rightarrow H_B^1((0, T); R^n)$  defines a homotopy. It is clear that the fixed points of  $H_\lambda$  are precisely the solutions to (2.1) $_\lambda$ . Now  $H_\lambda$  is fixed point free on  $\partial V$ . Moreover, the complete continuity of  $j$  together with the continuity of  $N^{-1}$  and  $F_\lambda$  imply that the homotopy  $H_\lambda$  is compact. Now  $H_0$  is essential so Theorem 1.5 of [9] implies that  $H_1$  is essential. Thus (2.1) has a solution. ■

We are now in a position to prove our basic existence theorem for first order initial problems.

**Theorem 2.3.** Suppose  $f: [0, T] \times R^n \rightarrow R^n$  satisfied (2.2) and (2.3). Then the initial value problem (2.1) has a solution in  $H^1((0, T); R^n)$  for

$$T < \int_0^\infty \frac{dx}{\psi(x)}.$$

Proof: To prove existence of a solution  $H^1((0, T); R^n)$  we apply Theorem 2.2. So to establish a priori bounds for (2.1) $_\lambda$ , let  $y(t)$  be a solution to (2.1) $_\lambda$ . Then since

$$|\lambda f(t; y)| \leq \psi(|y|)$$

we have for each  $t \in [0, T]$

$$\int_0^t \frac{|y'(u)|}{\psi(|y(u)|)} du \leq t \leq T.$$

Suppose  $y(t) \neq 0$  for some  $t \in [0, T]$ . Since  $y(0) = 0$  there is an interval  $[a, t]$  in  $[0, T]$  such that  $|y(s)| > 0$  on  $a < s \leq t$  and  $y(a) = 0$ . Then for any point  $t$  where  $y(t) \neq 0$

$$\int_a^t |y(x)|' ds = \int_a^t \frac{y(s) \cdot y'(s)}{|y(s)|} \leq \int_a^t |y'(s)| ds$$

and this together with the above inequality yields

$$\int_a^t \frac{|y(s)|'}{\psi(|y(s)|)} ds \leq T < \int_0^\infty \frac{dx}{\psi(x)}.$$

Theorem 1.2 yields

$$\int_0^{|y(t)|} \frac{dx}{\psi(x)} \leq T < \int_0^\infty \frac{dx}{\psi(x)}.$$

So there exists a constant  $M < \infty$  such that  $|y(t)| < M$ . Moreover,

$$\|y\|_{L^2} = \left( \int_0^T |y(t)|^2 dt \right)^{\frac{1}{2}} \leq MT^{\frac{1}{2}} \equiv M_1.$$

Also (2.3) and the differential equation yield

$$\left( \int_0^T |y'(t)|^2 dt \right)^{\frac{1}{2}} \leq \left( \int_0^T [\psi(|y(t)|)]^2 dt \right)^{\frac{1}{2}} \leq M_2 T^{\frac{1}{2}}$$

where  $M_2 = \text{ess sup}_{x \leq M_1} \psi(x)$ . Hence  $\|y\|_{H^1} \leq K = M_1 + M_2 T^{\frac{1}{2}}$  and the existence of a solution to (2.1) is established. ■

**Theorem 2.4 (Maximal Interval of Existence).** *Let  $\psi: [0, \infty) \rightarrow (0, \infty)$  be continuous. Then the initial value problem*

$$\begin{cases} y' = \tilde{f}(t; y), & t \in [0, \hat{T}] \\ y(0) = 0 \end{cases} \quad (2.4)$$

where  $\tilde{f}(t; y) = (\psi(|y|), 0, \dots, 0)$ , has a solution if and only if

$$\hat{T} < T_\infty = \int_0^\infty \frac{dx}{\psi(x)}.$$

**Proof:** The existence of a solution for  $\hat{T} < T_\infty$  follows from Theorem 2.3. Finally,  $y(t) = (y_1(t), \dots, y_n(t))$  is a solution to (2.4) if and only if  $y_2(t) = \dots = y_n(t) \equiv 0$  and  $y_1'(t) = \psi(|y_1|)$ ,  $y_1(t) > 0$ . Clearly,  $y_1' > 0$  and thus  $y_1(t) \geq 0$  since  $y_1(0) = 0$ . Now integration yields

$$\hat{T} = \int_0^{\hat{T}} \frac{y_1'(t)}{\psi(y_1(t))} dt = \int_0^{y_1(\hat{T})} \frac{dx}{\psi(x)} < \int_0^\infty \frac{dx}{\psi(x)} = T_\infty. \quad \blacksquare$$

Theorem 2.2 also holds for the inhomogeneous initial condition  $y(0) = \tau$ , via ideas of Theorem 5.1 in [10]. So trivial adjustments in the proof of Theorem 2.3 yields:

**Theorem 2.5.** Suppose  $f: [0, T] \times R^n \rightarrow R^n$  satisfies (2.2) and (2.3). Then the initial value problem

$$\begin{cases} y' = f(t; y), & t \in [0, T] \\ y(0) = r \end{cases} \quad (2.5)$$

has a solution in  $H^1((0, T); R^n)$  for each

$$T < \int_{|r|}^{\infty} \frac{dx}{\psi(x)}.$$

Moreover, this result is best possible as described in Theorem 2.4.

Example 1: (Electrical Circuits) Let us consider an electrical circuit which contains a resistance  $R$ , a condenser of capacitance  $C$ , and switch  $S$  and a generator  $E$ . Suppose the switch is closed at  $t = 0$ . Then Kirchoff's, Ohm's, and Coulomb's Laws imply that the  $RC$  circuit satisfies

$$R \frac{dq}{dt} + \frac{1}{C} q(t) = E(t, q)$$

where  $q(t)$  denotes the charge on the capacitor at time  $t$  and  $E(t, q)$  the value of the voltage impressed on the circuit by  $E$ .

We also have  $q(0) = q_0$  where  $q_0$  is the charge on the capacitor at  $t = 0$ . Thus we are interested in the initial value problem

$$\begin{cases} \frac{dq}{dt} = -\frac{1}{RC} q + \frac{E(t, q)}{R} \\ q(0) = q_0. \end{cases} \quad (2.6)$$

Suppose  $|E(t, q)| \leq A(t)|q| + B(t)$  for bounded functions  $A(t), B(t) \geq 0$  and that  $E(t, q)$  satisfies the Caratheodory Conditions. Consequently, since

$$\int_{|q_0|}^{\infty} \frac{dq}{A_0 q + B_0} = \infty,$$

for any constants  $A_0, B_0 > 0$ , Theorem 2.5 implies that (2.5) has a solution in  $H^1((0, T); R^n)$  for any  $T > 0$ . On the other hand suppose

$$|E(t, q)| \leq A(t)|q|^m + B(t), \quad m = 1, 2, \dots$$

for bounded functions  $A(t), B(t), \geq 0$  and that  $E(t, q)$  satisfies the Caratheodory Conditions. Then Theorem 2.5 implies that (2.5) has a solution in  $H^1((0, T); R)$  for any  $T < T_{\infty}$  where  $T_{\infty}$  is as described in Example 2 of [12].

### Acknowledgements.

We would like to thank Professor's Andrzej Granas and John W. Lee for their invaluable assistance during the research and writing up of this paper.

### References

1. H. Amann and M.G. Crandall, *On some existence theorems for semilinear elliptic equations*, Indiana Univ. Math. J. **27** (1978), 779-790.
2. H. Brezis, "Analyse Fonctionnelle, Theorie et Applications", Masson, Paris, 1983.
3. K.C. Chang, *The obstacle problem and partial differential equations with discontinuous nonlinearities*, Comm. Pure and App. Math. **33** (1980), 117-146.
4. J. Dugundji and A. Granas, "Fixed Point Theory", Vol. 1, Monographie Matematyczne, PNW, Warszawa, 1982.
5. S. Fücik, "Soluability of Nonlinear Equations and Boundary Value Problems", D. Reidel Pub. Co., Dordrecht-Boston and London, 1980.
6. S. Fücik and A. Kufner, "Nonlinear Differential Equations", Elsevier Scientific Pub. Co, Amsterdam-Oxford and New York, 1980.
7. A. Granas, *Sur La Methode de Continuité de Poincaré*, Comptes Rendus Acad.Sci. **282** (1976), 983-985, Paris.
8. A. Granas, R.B. Guenther and J.W. Lee, *Nonlinear boundary value problems for classes of ordinary differential equations*, Rocky Mountain J. of Math. **10** (1980), 35-58.
9. A. Granas, R.B. Guenther and J.W. Lee, *Topological transversality I (some nonlinear diffusion problems)*, Pac.Jour.Math. **89** (1980), 53-67.
10. R.B. Guenther, "Problems Aux Limites Non Lineaires Pour Certaines Classes D'Equations Differentielles Ordinaires", Les Presses de L'Université Montreal, 1985.
11. M.A. Krasnosel'skii, "Topological Methods in the Theory of Nonlinear Integral Equations", MacMillan Company, New York, 1964.
12. J.W. Lee and D. O'Regan, *Topological transversality IV, applications to initial value problems*, Annales Polonici Mathematici **48**.
13. H.L. Royden, "Real Analysis", 3rd edition, MacMillan Pub. Co., New York, 1988.
14. W. Rudin, "Real and Complex Analysis", McGraw-Hill, New York, 1966.

15. M.M. Vainberg, "Variational Methods for the Study of Nonlinear Operators", Holden-Day Inc., San Francisco, London and Amsterdam, 1964.
16. F.-C. Liu and M. Frigon, *Remarque sur une application de l'indicatrice de Banach au changement de variable dans une intégrale*, Rapport de Recherche de l'université de Montréal (to appear).
17. M. Frigon, *Méthode de transversalité topologique appliquée à des Problèmes nonlineares pour des équations différentielles ordinaires*, Ph.D. Thèse, Université de Montréal.